

Results on High p_T Particle Production from the PHENIX Experiment at RHIC

K. Reygers^a for the PHENIX Collaboration

^aUniversity of Münster, D-48149 Münster, Germany

Transverse momentum (p_T) spectra of neutral pions and charged hadrons measured in Au+Au and d+Au collisions at $\sqrt{s_{NN}} = 200$ GeV by the PHENIX experiment at RHIC are compared to p+p reference spectra at the same $\sqrt{s_{NN}}$. In central Au+Au collisions a factor 4 – 5 suppression for neutral pions and charged hadrons with $p_T > 5$ GeV/ c is found relative to the p+p reference scaled by the nuclear overlap function $\langle T_{AB} \rangle$. In contrast, such a suppression of high p_T particles is absent in d+Au collisions.

1. Introduction

The objective of the heavy ion program at the Relativistic Heavy Ion Collider (RHIC) at the Brookhaven National Laboratory is to create and study a deconfined and thermalized state of strongly interacting matter, the quark-gluon plasma. In a central Au+Au collision at RHIC with a center-of-mass energy per nucleon-nucleon pair of $\sqrt{s_{NN}} = 200$ GeV about 5000 charged particles are produced. The bulk of the particles carry rather low transverse momenta ($p_T < 2$ GeV/ c) with respect to the beam axis. Of particular importance for the study of nucleus-nucleus (A+A) collisions, however, is the small fraction of particles with higher transverse momentum. These particles are expected to be produced in parton-parton interactions with high momentum transfer (*hard processes*). In proton-proton interactions the scattered partons propagate through the normal QCD vacuum where they finally fragment into the observed, colorless hadrons. In A+A collisions the hard parton-parton scatterings take place in the early stage of the collision, before a possible quark-gluon plasma has formed. The highly-energetic scattered partons traverse the subsequently produced excited nuclear matter. As a consequence, particle production at high transverse momentum is sensitive to properties of the hot and dense matter created in A+A collisions.

2. High p_T Particle Yields in p+p, d+Au and Au+Au

Neutral pions and charged particles at mid-rapidity were measured with the two central spectrometer arms of the PHENIX experiment. Each arm covers $|\eta| \leq 0.35$ in pseudorapidity and $\Delta\phi = \pi/2$ in azimuth. Neutral pions were measured by the PHENIX electromagnetic calorimeters (EMCal) via the $\pi^0 \rightarrow \gamma\gamma$ decay. The EMCal consists of six lead-scintillator and two lead-glass sectors, each located at a radial distance of about 5 m to the interaction region. The lateral segmentation of the EMCal sectors is




Figure 1. Neutral pion invariant yields in p+p and central (0 – 10% of $\sigma_{\text{geo}}^{\text{AuAu}}$) Au+Au collisions at $\sqrt{s_{\text{NN}}} = 200$ GeV and in central (0 – 10.8% of $\sigma_{\text{geo}}^{\text{PbPb}}$) Pb+Pb collisions at $\sqrt{s_{\text{NN}}} = 17.3$ GeV. The p+p spectra are scaled by the number of inelastic nucleon-nucleon collisions for the respective A+A reaction ($\langle N_{\text{coll}}^{\text{Au+Au}} \rangle = 955 \pm 94$, $\langle N_{\text{coll}}^{\text{Pb+Pb}} \rangle = 651 \pm 65$). For the p+p π^0 spectrum at $\sqrt{s} = 17.3$ GeV a parameterization from [4] was used.

$\Delta\eta \times \Delta\phi \approx 0.01 \times 0.01$. The systematic uncertainty of the absolute energy scale of the EMCal is less than 1.5%. The charged particle spectrometer consists of a magnetic field which is axially symmetric around the beam axis and tracking detectors. Charged particle tracks are reconstructed with a drift chamber followed by two layers by multiwire proportional chambers with pad readout. The absolute momentum scale is known to 0.7%.

For hard parton-parton interactions with small cross section a A+A collision can be considered as a incoherent sequence of individual nucleon-nucleon collisions. Thus, in the absence of nuclear effects the yield of particles produced in hard processes is expected to scale with the average number of inelastic nucleon-nucleon collisions $\langle N_{\text{coll}} \rangle$. This picture of a incoherent superposition of nucleon-nucleon or parton-parton interactions breaks down for processes which yield low- p_T particles ($p_T < 1 - 2$ GeV/c). Particle production in this p_T range is known to scale approximately with the number of participants N_{part} , *i.e.* with the number of nucleons which underwent at least one inelastic nucleon-nucleon collision. In order to quantify nuclear medium effects on high p_T particle production, a nuclear modification factor is defined as

$$R_{\text{AB}}(p_T) = \frac{(1/N_{\text{evt}}^{\text{AB}}) d^2 N_{\text{AB}} / dp_T d\eta}{\langle T_{\text{AB}} \rangle \times d^2 \sigma_{\text{pp}} / dp_T d\eta} = \frac{(1/N_{\text{evt}}^{\text{AB}}) d^2 N_{\text{AB}} / dp_T d\eta}{\langle N_{\text{coll}} \rangle \times (1/N_{\text{evt}}^{\text{pp}}) d^2 N_{\text{pp}} / dp_T d\eta}. \quad (1)$$

The average nuclear overlap function $\langle T_{\text{AB}} \rangle$ is determined solely from the geometry of the nuclei A and B and bears resemblance to the integrated luminosity of a particle collider because the average number of nucleon-nucleon collisions per A+B collision is given by $\langle N_{\text{coll}} \rangle = \sigma_{\text{inel}}^{\text{pp}} \times \langle T_{\text{AB}} \rangle$.

The invariant yields of neutral pions in p+p and central Au+Au collisions at $\sqrt{s_{\text{NN}}} =$




Figure 2. Nuclear modification factor $R_{AA}(p_T)$ for central Au+Au collisions at RHIC, central Pb+Pb collisions at CERN SPS, and $\alpha+\alpha$ collisions at the ISR.

200 GeV are shown in Figure 1 [2, 3]. The p+p spectrum is scaled by $\langle N_{\text{coll}} \rangle$. Compared to this reference a striking suppression of neutral pions is observed in central Au+Au collisions. The neutral pion spectrum for central Pb+Pb collisions at $\sqrt{s_{\text{NN}}} = 17.3$ GeV is also shown for comparison [4].

The nuclear modification factors determined from the spectra in Figure 1 are shown in Figure 2. For Pb+Pb at $\sqrt{s_{\text{NN}}} = 17.3$ GeV R_{AA} increases with p_T and exceeds the N_{coll} scaling expectation for $p_T > 2$ GeV/c. A similar observation was made in p+A reactions and has been called *Cronin-effect*. This effect is usually attributed to multiple soft nucleon-nucleon interactions which lead to a transverse momentum component of the parton-parton system before the hard scattering occurs. In view of the data from the CERN SPS the factor 4–5 neutral pion suppression observed in central Au+Au collisions at RHIC was a dramatic discovery.

Among the various theoretical explanations for the hadron suppression in central Au+Au collisions at RHIC is (i) energy loss of the scattered partons in the medium of high color charge density (mainly gluons) created in central Au+Au collisions [6], (ii) saturation of gluons at small Bjorken- x in the initial state wavefunction of the Au nuclei which leads to fewer hard gluon-gluon scatterings and thus to fewer high p_T particles [7], (iii) final state scattering of high p_T hadrons with other hadrons produced in the Au+Au collision [8]. Explanation (i) is consistent with the creation of a quark-gluon plasma whereas (ii) and (iii) explain the observed suppression without assuming a quark-gluon plasma. Explanations like (i) which attribute the high p_T hadron suppression in central Au+Au collision to final state effects and models which are based on initial state effects make different predictions for deuteron+gold collisions which were recently studied at RHIC. In a d+Au collision the highly excited matter is not created over a large volume and so high p_T particle suppression is not expected in the parton energy loss scenario (i). By contrast, in the framework of the gluon saturation model high p_T particle production ought to be suppressed in d+Au as well. The gluon saturation calculation in [9] predicts




Figure 3. Nuclear modification factor for neutral pions and charged hadrons in minimum bias d+Au collisions at $\sqrt{s_{NN}} = 200$ GeV. The results for Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV are shown for comparison.

$R_{AB} \approx 0.7 - 0.8$ for central d+Au collisions.

The experimental nuclear modification factors for neutral pions and charged hadrons in minimum bias d+Au collisions ($\langle N_{coll} \rangle \approx 8.5$) are shown in Figure 3. Neither high p_T neutral pion nor charged hadron yields are suppressed with respect to N_{coll} scaling. The measurements therefore disfavor models like the gluon saturation model which are based on initial state effects.

3. Conclusions

The suppression of high p_T particle production in central Au+Au collisions is one of the most interesting findings at RHIC. Parton energy loss, which is expected to occur in a quark-gluon plasma, is able to explain the measurements. In light of data from the d+Au beamtime at RHIC a competing theory, gluon saturation, appears as an unlikely explanation for the suppression in Au+Au.

REFERENCES

1. PHENIX collaboration, S.S. Adler et al., Phys. Rev. Lett. **91** (2003) 072303
2. PHENIX collaboration, S.S. Adler et al., Phys. Rev. Lett. **91** (2003) 072301
3. PHENIX collaboration, S.S. Adler et al., hep-ex/0304038
4. WA98 collaboration, M.M. Aggarwal et al, Eur. Phys. J. **C 23** (2002) 225
5. J.W. Cronin et al., Phys. Rev. **D11** (1975), 3105
6. M. Gyulassy, I. Vitev, X.N. Wang, B.W. Zhang, nucl-th/0302077
7. E. Iancu, R. Venugopalan, hep-th/0303204
8. K. Gallmeister, C. Greiner, Z. Xu, Phys. Rev. **C 67** (2003) 014008
9. D. Kharzeev, E. Levin, L. McLerran, Phys. Lett. **B 561** (2003), 93